

Comment on Dirac spectral sum rules for QCD₃

Shinsuke M. Nishigaki*

Department of Physics, Faculty of Science, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan
(October 2, 2000)

Recently Magnea [Phys. Rev. **D61**, 056005 (2000); Phys. Rev. **D62**, 016005 (2000)] claimed to have computed the first sum rules for Dirac operators in 3D gauge theories from 0D non-linear σ models. I point out that these computations are incorrect, and that they contradict with the exact results for the spectral densities unambiguously derived from random matrix theory by Nagao and myself.

Magnea [1] has recently claimed to have derived Dirac spectral sum rules for three-dimensional gauge theories coupled in a (P, \mathbf{Z}_2) -invariant manner to fundamental fermions with $N_c = 2$ (corresponding to the Dyson index $\beta = 1$) and adjoint fermions ($\beta = 4$). She employed the small-mass expansion of the low-energy effective theories, i.e. the zero-dimensional σ models over Riemannian symmetric spaces $\mathcal{M} = \text{CII}$ and BDI , instead of AIII that had been proposed for the case with fundamental fermions and $N_c \geq 3$ ($\beta = 2$) [2]. She concluded that the first sum rule in the presence of even number (N_f) of massless 2-component complex or 4-component real (Majorana) fermions is common both to the $\beta = 1$ and $\beta = 4$ universality classes, and takes the form[†]

$$\left\langle \sum_i \frac{1}{\zeta_i^2} \right\rangle^{(1,4)} = \int_{-\infty}^{\infty} d\zeta \frac{\rho_s^{(1,4)}(\zeta)}{\zeta^2} = \frac{4}{N_f}, \quad (1)$$

where ζ stands for an unfolded Dirac eigenvalue (i.e. rescaled by $1/(\pi\rho(0))$) and $\rho_s^{(\beta)}(\zeta)$ stands for the scaled

spectral density. If true, this conclusion, derived from an obviously correct formula (see Ref. [3])

$$\left\langle \sum_i \frac{1}{\zeta_i^2} \right\rangle = \frac{d^2}{N_f M} \quad (2)$$

(d stands for the rank of the matrix that parameterizes \mathcal{M} , and M for the dimension of \mathcal{M}), would be surprising, as the four-dimensional counterpart of the spectral sum rules are known to be different for three values of β [4,3].

On the other hand, Nagao and myself [5] have obtained Pfaffian expressions for the generic p -level correlation functions in a presence of an arbitrary number of (for $\beta = 1$) and an arbitrary number of pairwise degenerate (for $\beta = 4$) finite fermion masses $\{\mu_f\}$, by applying the skew-orthogonal polynomial method to pertinent random matrix ensembles. To make comparison with Eq.(1), I take a completely confluent limit $\mu_f \rightarrow 0$ for all f , of our results [Ref. [5], Eqs.(2.40), (2.42), (3.23), (3.25), (3.49), (3.51)] with $p = 1$ (spectral density), to obtain:

$$\pi \rho_s^{(1)}(\zeta) = 1 - \frac{3}{\zeta^2} + \frac{3}{2\zeta^4} - \frac{3 \cos 2\zeta}{2\zeta^4} \quad (N_f = 2), \quad (3a)$$

$$= 1 - \frac{10}{\zeta^2} - \frac{30}{\zeta^4} + \frac{210}{\zeta^6} + \frac{525}{2\zeta^8} + \left(\frac{70}{\zeta^5} - \frac{525}{\zeta^7} \right) \sin 2\zeta + \left(-\frac{5}{\zeta^4} + \frac{315}{\zeta^6} - \frac{525}{2\zeta^8} \right) \cos 2\zeta \quad (N_f = 4), \quad (3b)$$

$$= 1 - \frac{21}{\zeta^2} - \frac{357}{2\zeta^4} - \frac{945}{\zeta^6} + \frac{48195}{\zeta^8} + \frac{218295}{\zeta^{10}} + \frac{1964655}{2\zeta^{12}} + \left(\frac{378}{\zeta^5} - \frac{50652}{\zeta^7} + \frac{873180}{\zeta^9} - \frac{1964655}{\zeta^{11}} \right) \sin 2\zeta \\ + \left(-\frac{21}{2\zeta^4} + \frac{5859}{\zeta^6} - \frac{266490}{\zeta^8} + \frac{1746360}{\zeta^{10}} - \frac{1964655}{2\zeta^{12}} \right) \cos 2\zeta \quad (N_f = 6), \quad (3c)$$

$$\pi \rho_s^{(4)}(\zeta) = 1 - \frac{\sin 2\zeta}{2\zeta} \quad (N_f = 2), \quad (4a)$$

$$= 1 - \frac{\sin^2 2\zeta}{4\zeta^2} + \left(-\frac{\sin 2\zeta}{4\zeta^2} + \frac{\cos 2\zeta}{2\zeta} \right) \text{Si}(2\zeta) \quad (N_f = 4), \quad (4b)$$

$$= 1 - \frac{3}{4\zeta^2} + \frac{3}{32\zeta^4} + \left(\frac{1}{\zeta} - \frac{3}{4\zeta^3} \right) \sin 2\zeta + \frac{3 \cos 2\zeta}{2\zeta^2} - \frac{3 \cos 4\zeta}{32\zeta^4} \quad (N_f = 6), \quad (4c)$$

$$= 1 - \frac{27}{16\zeta^2} + \frac{45}{64\zeta^4} + \frac{45}{256\zeta^6} + \left(\frac{15}{32\zeta^3} - \frac{45}{64\zeta^5} \right) \sin 4\zeta + \left(-\frac{3}{16\zeta^2} + \frac{45}{64\zeta^4} - \frac{45}{256\zeta^6} \right) \cos 4\zeta \\ + \left[\left(\frac{9}{4\zeta^2} - \frac{45}{32\zeta^4} \right) \sin 2\zeta + \left(-\frac{3}{4\zeta} + \frac{45}{16\zeta^3} \right) \cos 2\zeta \right] \text{Si}(2\zeta) \quad (N_f = 8), \quad (4d)$$

(Si stands for the sine-integral function) and so forth. These expressions for the spectral densities lead to the sum rules

$$\int_{-\infty}^{\infty} d\zeta \frac{\rho_s^{(1)}(\zeta)}{\zeta^2} = \frac{N_f}{(N_f - 1)(2N_f + 1)}, \quad (5)$$

$$\int_{-\infty}^{\infty} d\zeta \frac{\rho_s^{(4)}(\zeta)}{\zeta^2} = \frac{N_f}{(N_f - 1)(N_f/2 + 1)}, \quad (6)$$

which are sensitive to the Dyson index β . The above sum rules agree perfectly with the numerical results for random matrix ensembles of large but finite ranks (~ 40), obtained by Hilmoine and Niclasen [6] via two alternative methods (an analytical method of Widom's [Table 4 of Ref. [6]] and numerical Monte-Carlo simulations of random matrix ensembles). Therefore I conclude that the expression (1) is erroneous, and the coincidence of the first sum rules for $\beta = 1$ and $\beta = 4$ claimed in her papers is illusory.

I thank C. Hilmoine for bringing this issue to my attention. This work was supported in part by JSPS, and by

Grant-in-Aid No. 411044 from the Ministry of Education, Science and Culture, Japan.

* e-mail address: nishigak@th.phys.titech.ac.jp

† N_f in Ref. [1] stands for the number of 4-component spinors, i.e. a half of N_f in this note and in Refs. [2,5,6].

- [1] U. Magnea, Phys. Rev. **D61**, 056005 (2000); Phys. Rev. **D62**, 016005 (2000).
- [2] J.J.M. Verbaarschot and I. Zahed, Phys. Rev. Lett. **73**, 2288 (1994).
- [3] A. Smilga and J.J.M. Verbaarschot, Phys. Rev. **D51**, 829 (1995); M.A. Halasz and J.J.M. Verbaarschot, Phys. Rev. **D52**, 2563 (1995).
- [4] H. Leutwyler and A. Smilga, Phys. Rev. **D46**, 5607 (1992).
- [5] T. Nagao and S.M. Nishigaki, hep-th/0005077, Phys. Rev. **D62** (2000) in press.
- [6] C. Hilmoine and R. Niclasen, hep-th/0004081.